

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/57171 A1

(51) International Patent Classification⁷: **C11D 3/37**,
1/62, 3/00

(21) International Application Number: **PCT/EP01/00409**

(22) International Filing Date: 15 January 2001 (15.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00400282.0 2 February 2000 (02.02.2000) EP

(71) Applicant (for AE, AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, KE, LC, LK, LS, MN, MW, NZ, SD, SG, SI, SZ, TT, TZ, UG, ZA, ZW only): **UNILEVER PLC** [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).

(71) Applicant (for all designated States except AE, AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, IN, KE, LC, LK, LS, MN, MW, NZ, SD, SG, SI, SZ, TT, TZ, UG, ZA, ZW): **UNILEVER NV** [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).

(71) Applicant (for IN only): **HINDUSTAN LEVER LIMITED** [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Maharashtra, 400 020 Mumbai (IN).

(72) Inventors: **AUBAY, Eric**; 1 ter rue de Metz, F-94170 Le Perreux sur Marne (FR). **BLOKZIJL, Wilfried**; Unilever Research Port Sunlight, Quarry Road East, Bebington, Wirral, Merseyside CH63 3JW (GB). **GEFFROY, Cedric**; 18 rue Chaptal, F-75009 Paris (FR). **MAHMOOD, Khalid**; Unilever Research Port Sunlight, Quarry Road East, Bebington, Wirral, Merseyside CH63 3JW (GB).

THOMPSON, Laurence, Griffith; Unilever Research Port Sunlight, Quarry Road East, Bebington, Wirral, Merseyside CH63 3JW (GB). **WALSH, John, Michael**; Unilever Research Port Sunlight, Quarry Road East, Bebington, Wirral, Merseyside CH63 3JW (GB). **WONG, Kenneth**; 41 rue du Faubourg St Denis, F-75010 Paris (FR). **YEUNG, Dominic**; 3661 Golden Orchard Drive, Mississauga, Ontario L4Y 3J2 (CA).

(74) Agents: **FRANSELLA, Mary, Evelyn** et al.; Unilever PLC, Patent Department, Colworth House, Sharnbrook, Bedford, Bedfordshire MK44 1LQ (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report
before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/57171 A1

(54) Title: **POLYMERS FOR LAUNDRY APPLICATIONS**

(57) Abstract: A polymer for use in a composition or method of laundry treatment is a copolymer of: (a) one or more anionic monomer units; (b) one or more cationic monomer units; and (c) optionally, one or more uncharged monomer units; wherein, the number ratio of the total of all negative charges on the anionic monomer unit(s) to the total of all positive charges on the cationic monomer unit(s) is from 10 : 1 to 3 : 1, especially from 17 : 3 to 3 : 1.

POLYMERS FOR LAUNDRY APPLICATIONS

5

Field of Invention

The present invention relates to polymers which are used in laundry cleaning products, e.g. for incorporation in products for dosing in the wash and/or rinse. They are intended
10 for, but not limited to, soil release benefits in such products.

Background of the Invention

15 The term "soil release polymer" is used in the art to cover polymeric materials which assist release of soil from fabrics, e.g. cotton or polyester based fabrics. For example, it is used in relation to polymers which assist release of soil direct from fibres. It is also used to refer to polymers which modify the fibres so that dirt adheres to the polymer-modified fibres rather than to the fibre material itself. Then, when the fabric is washed
20 the next time, the dirt is more easily removed than if it was adhering the fibres. Although not wishing to be bound by any particular theory or explanation, the inventors believe that the soil release polymers utilised in the present invention probably exert their effect mainly by the latter mechanism.

25 WO-A-98/23714 discloses water-soluble use of nitrogen-containing soil-release polymers in detergent products. Some of these polymers are formed from acrylamide monomers polymerised with co-monomers which are amines of alkylacrylates. These materials are essentially neutral, i.e. have no substantial overall positive or negative charge. Thus, they are suited to soil release from polyester rather than from cotton.

30

Another type of nitrogen-containing soil release polymer described in WO-A-98/23714 is formed from dicarboxylic acid monomers and hydrophilic co-monomers which are secondary amines which contain (poly)alkyleneoxy groups. Since these molecules do not have an overall net positive charge, they are also non-substantive to cotton.

5

A third type of nitrogen-containing soil-release polymer disclosed in WO-A-98/23714 is formed from alkyleneacrylate monomers having a terminal quaternary ammonium group and co-monomers which are (meth)acrylic acid or esters or salts thereof.

Therefore, such a polymer may have both cationic and anionic groups. Only a single example of such a material is given. This is polymer formed from a cationic monomer, namely methacrylamidopropyl trimethylammonium chloride (MAPTAC), and anionic monomer, namely acrylic acid (AA) and a neutral monomer, namely isobutylacrylate (IsoBuA). These particular cationic and anionic monomers carry, respectively, a single positive and negative charge. However, this particular kind of polymer has proved to be not very suited to soil release.

10
15

US-A-3 749 682 discloses copolymers of polyvinylpyrrolidone (PVP) and vinyl acetate for use as soil anti-redeposition agents.

WO-A-97/42285 discloses cotton soil-release polymers comprising a polyamine backbone and quaternary ammonium cationic groups. However, the present applicants have found that whilst cationic groups give good substantivity to cotton, a high density of cationic positive charges on the polymer gives rise to staining of the fabric.

EP-A-0995791 discloses a broad range of hydrophobically modified polycarboxylate polymers which are said to be useful for promoting soil release from fabrics, particularly cotton and cotton-containing fabrics. The polymers can comprise up to three moieties A, B and C wherein A is a polymerised residue of a monomer selected from one or more C₃-C₈ monoethylenically unsaturated carboxylic acids, B is a polymerised residue of a monomer selected from one or more C₃-C₆₀ alkyl (meth)acrylates, ethoxylated C₁-C₂₄ alkyl (meth)acrylates, and poly(alkylene glycol)

25
30

- (meth)acrylates, alkyl or aromatic ethers of poly(alkylene glycol) and the corresponding maleate mono and di-esters thereof, and C is a polymerised residue of a monomer selected from one or more ethylenically unsaturated monomers which are copolymerisable with the monomers in A and B. However, only two specific examples are given which contain both anionic and cationic groups. One such polymer is formed from an anionic monomer, namely acrylic acid (AA), a cationic monomer, namely dialkyldimethyl ammonium chloride (DADMAC), and lauryl (C₁₂) methacrylate ethoxylated with 4 moles of ethylene oxide (E4LMA) whereas the other polymer is formed from AA, E4LMA and [2-(methacryloyloxy)ethyl]trimethyl ammonium chloride (MAETMAC) as a cationic monomer. These particular anionic and cationic monomers carry, respectively, a single negative and positive charge. However, it is clear from the examples in question that the number ratio of anionic : cationic charges is >50:1.
- 15 According to GB-A-2 104 091, copolymers of anionic and cationic vinyl monomers can be used as detergency builders, for a wide range of detergent products, e.g. for warewashing, hard surface cleaning, textile cleaning and hair products. However, there is no disclosure that these materials are capable of acting as soil-release polymers. They are not well suited to soil release from cotton and can lead to significant staining.
- 20 Although a wide range of anionic : cationic mole ratios is claimed (from 1 : 99 to 99 : 1), in all of the examples, the number ratio of anionic (negative) : cationic (positive) charges is 1 : 1, which again, is not suited to providing the cotton substantivity required of a cotton soil-release polymer.
- 25 US-A-5783533 discloses various amphoteric copolymers as rheological modifiers of lamellar phases of detergent or cosmetic compositions. However, there is no disclosure that these materials are capable of acting as soil-release polymers.
- JP-A-59135293 discloses detergent compositions which contain an amphoteric copolymer consisting of at least 10 mol% cationic vinyl monomer units, at least 10 mol% anionic vinyl monomer units and at least 10 mol% nonionic vinyl monomer units.

The preferred proportion of cationic vinyl monomer units to anionic vinyl monomer units by molar ratio from 1:2 to 2:1. However, there is no disclosure that these materials are capable of acting as soil-release polymers.

- 5 US-A-5413731 discloses water-soluble terpolymers which are useful in automatic machine dishwashing detergent formulations. These terpolymers contain as polymerised units (a) from about 92 to about 30% by weight of one or more C₃-C₆ monoethylenically unsaturated carboxylic acids, (b) from about 5 to about 50% by weight of one or more aminoacryloyl derivatives, and (c) from about 25% by weight of one or more monoethylenically unsaturated monomers polymerisable with (a) and (b).
10 However, there is no disclosure that these materials could be utilised in a composition for washing and/or rinsing landry, nor any suggestions that these terpolymers could act as soil-release polymers.
- 15 Thus, there remains a need to obtain soil release in laundry products based on soil-release agents which are copolymers of cationic and anionic monomers, give better substantivity to cotton and therefore, better soil release from cotton fabrics, fewer or no detrimental interactions with LAS or other anionic surfactants, as well as less staining negatives.

20

Definition of the Invention

- A composition for washing and/or rinsing of laundry, the composition comprising one or more surfactants suitable for use in laundry wash and/or rinsing products and a polymer which is a co-polymer formed of:-
25

- (a) one or more anionic monomer units;
- 30 (b) one or more cationic monomer units; and

(c) optionally, one or more neutral (uncharged) monomer units;

wherein, the number ratio of the total of all negative charges on the anionic monomer unit(s) to the total of all positive charges on the cationic monomer unit(s) is from 10 : 1 to 3 : 1, preferably from 8 : 1 to 3 : 1, especially from 17 : 3 to 3 : 1.

The applicants have found that these charge ratios (progressively) give the optimum balance of soil release performance with avoidance of staining and detrimental interactions with the LAS or other anionic surfactants.

10

A second aspect of the invention provides use of a polymer for a method effecting soil-release of a laundry item, the method comprising contacting the laundry item with the said polymer, the polymer being a copolymer formed of:-

15 (a) one or more anionic monomer units;

(b) one or more cationic monomer units; and

(c) optionally, one or more uncharged monomer units;

20

wherein, the number ratio of the total of all negative charges on the anionic monomer unit(s) to the total of all positive charges on the cationic monomer unit(s) is from 10 : 1 to 3 : 1, preferably from 8 : 1 to 3 : 1, especially from 17 : 3 to 3 : 1.

25 This aspect of the invention may also be expressed as a process for effecting soil release from a laundry item, said process comprising contacting the laundry item with the said polymer, the polymer being a copolymer formed of:-

(a) one or more anionic monomer units;

30

(b) one or more cationic monomer units; and

(c) optionally, one or more uncharged monomer units;

wherein, the number ratio of the total of all negative charges on the anionic monomer unit(s) to the total of all positive charges on the cationic monomer unit(s) is from 10 : 1 to 3 : 1, preferably from 8 : 1 to 3 : 1, especially from 17 : 3 to 3 : 1.


Detailed Description of the Invention

The Polymer

The polymer is a co-polymer containing one or more anionic monomer units, one or more cationic monomer units and optionally, one or more uncharged monomer units.


Preferably, the polymer has a weight average molecular weight of from 500 or from 1,000 to 1,000,000 or to 500,000, more preferably from 50,000 to 250,000 as determined by the methods of viscosity measurement and GPC using a polyethyleneoxide as standard.

The anionic monomer unit(s) (a) is/are preferably selected from one or more units derived from ethylenically unsaturated monomers having at least one anionic group. Typical such monomers have the general formula (A)



wherein at least two of Q^1 - Q^4 are independently selected from hydrogen and methyl;

either one or two of Q^1 - Q^4 are independently selected from anionic groups, preferably of formula:



- 5 wherein either or both of Q^5 and Q^6 is/are absent, Q^5 otherwise representing -Ph-, -CO-, -CH₂=CH₂, -CONH- or -CO-O- and Q^6 otherwise representing a C₁₋₄ alkylene linkage, one or more of the hydrogen atoms of which is independently optionally substituted by an -OH group or a group -Y;
- 10 Y is selected from groups of formula -CO₂H, -SO₃H, -OSO₃H, -PO₄H, -PO₃H, -OPO₃H₂ and -OPO₃H₃;

- and in the case where two only of Q^1 - Q^4 are independently hydrogen or methyl and only one of Q^1 - Q^4 is - Q^5 - Q^6 -Y, then the remaining group of Q^1 - Q^4 can be any other
- 15 compatible uncharged group, for example aliphatic, aromatic or mixed aliphatic-aromatic groups having from 2 to 20 carbon atoms (optionally also containing one or more heteroatoms) such as C₂₋₂₀ alkyl groups, C₅₋₁₂ cycloalkyl groups, C₅₋₉ aryl groups, C₁₋₈ alkyl-C₅₋₉ aryl groups, any cycloalkyl or aryl group optionally containing one or two heteroatoms independently selected from nitrogen, oxygen and sulphur.

20

- Preferred anionic groups for the anionic monomer units (whether or not derived from monomers of formula (A)) are selected from -CO₂H, -SO₃H, -OSO₃H, -CH₂OSO₃H, -CH=CHSO₃H and groups of formula -(CO)_p-CH₂-CQ⁷Q⁸CO₂H, -PO₄H, -PO₃H, -OPO₃H₂, -OPO₃H₃, wherein p is 0 or 1, Q⁷ is selected from H and OH and Q⁸ is
- 25 selected from H and CO₂H; and salts thereof.


- A non-limiting list of suitable ethylenically unsaturated anionic monomers includes acrylic acid, methacrylic acid, α -ethacrylic acid, β , β -dimethylacrylic acid, methylenemalonic acid, vinylacetic acid, allylacetic acid, ethylideneacetic acid,
- 30 propylideneacetic acid, crotonic acid, maleic acid or anhydride, fumaric acid, itaconic

acid, citraconic acid, mesaconic acid, N-(methacryloyl)alanine, N-(acryloyl)-hydroxyglycine, sulphopropyl acrylate, sulphoethyl acrylate, sulphoethyl methacrylate, styrenesulphonic acid, vinylsulphonic acid, 2-sulphoethyl methacrylate, sodium allyloxy hydroxypropyl sulphonate, vinylphosphonic acid, phosphoethyl acrylate, phosphonoethyl acrylate, phosphopropyl acrylate, phosphonopropyl acrylate, phosphoethyl methacrylate, phosphonoethyl methacrylate, phosphopropyl methacrylate, phosphonopropyl methacrylate, ethyleneglycol methacrylate phosphate, sulphate of alkoxylate (meth)acrylate, and salts thereof.

Any reference herein to an alkyl group on its own or as part of another group includes reference to straight and branched forms thereof.


Any anionic group forming part of an anionic monomer starting material or anionic monomer unit of the polymer may be in the acid form or salt form. Often, the free acid form may be neutralised either as part of the process for forming the polymer or when the polymer is incorporated in the detergent composition. Suitable counter-cations of the salt forms are alkali metals such as sodium or potassium, alkaline earth metals such as magnesium or organic ions such as NH_4^+ .

The monomer unit(s) (b) are preferably derived from ethylenically unsaturated monomers and advantageously comprise at least one quaternary ammonium group. Preferably they are selected from the units derived from compounds of following general formulae (I) to (III):



in which:

- R^1 is a hydrogen atom or a methyl group, preferably a methyl group;
- R^2 , R^3 and R^4 are linear or branched $C_1 - C_6$ alkyl groups;
- n is from 1 to 4, in particular the number 3;
- Z^1 is a group $-C(O)O$, $-C(O)NH-$ or $-O-$; and
- 5 - X^- is a counterion compatible with the water-soluble nature of the polymer;




10

in which:

- R^5 and R^8 are, independently hydrogen, or a linear or branched $C_1 - C_6$ alkyl group;
- 15 - R^6 and R^7 are independently represent alkyl, hydroxyalkyl or aminoalkyl group in which the alkyl group is a linear or branched $C_1 - C_6$ chain, preferably a methyl group;
- m and p are independently from 1 to 3; and
- X^- is as defined in formula (I); and

20



in which:


- 25 - R^9 is hydrogen, methyl or ethyl;

- R^{10} , R^{11} , R^{12} , R^{13} and R^{14} independently selected from groups as defined for R^6 and R^7 in formula(II);
- q is from 0 to 10, preferably from 0 to 2;
- r is from 0 to 6, preferably from 1 to 6, more preferably from 2 to 4;
- 5 - Z^1 is as defined in formula (I);
- Z^2 represents a $(CH_2)_s$ group, s being from 1 to 6, preferably from 2 to 4;
- Z^3 is a linear or branched $C_2 - C_{12}$, advantageously $C_3 - C_6$, polymethylene chain optionally interrupted by one or more heteroatoms or heterogroups, in particular O or NH, and optionally substituted by one or more hydroxyl or amino groups, preferably hydroxyl groups; and
- 10 - X^- , is as defined in formula (I); and

and also from ethylenically unsaturated monomers containing an aliphatic or aromatic cyclic moiety which contains a charged nitrogen (N^+) atom.

15


Preferred monomers of formula (I) are those of (IV):



- 20 wherein Z^1 is as defined in formula(I), especially that where Z^1 is -NH- and the counterion is a chloride (MAPTAC).

A preferred monomer of formula (II), is:-


25



in which X is a chloride (DADMAC).

5 Preferred monomers of formula (III) are those wherein:

- q is 2 or 3, especially 3;
- r is from 0 to 2, more preferably 0 to 1, especially 0;
- Z³ is




10


where t is from 1 to 4, preferably 1, and R¹⁰ to R¹⁴ which are the same or different, and represent a methyl or ethyl group.

Particularly preferred monomers (b) of the latter type are those of following formula:

15



wherein r is from 2 to 4, and more particularly the monomer




X- representing the chloride ion (Diquat)

- 5 When one or more uncharged monomer units (c) are also incorporated in the polymer, they are in a mole ratio of total uncharged units relative to the total of all anionic monomer units plus the total of all cationic monomer units of from 99:1 to 0:100, more preferably from 5:1 to 0:1. In another preferred embodiment, the mole ratio of the uncharged units relative to the total of all anionic monomer units plus the total of all
- 10 cationic monomer units is from 1 : 1 to 1 : 99, preferably from 3 : 2 to 1 : 9.

Preferably, the uncharged monomer units are derived from ethylenically unsaturated monomers, suitably selected from one or more of the following types (i)-(iii):-

- 15
- (i) hydrophilic neutral monomers such as (meth)acrylamide and their N-monosubstituted or N,N-disubstituted versions (such as N-isopropylacrylamide, N-butylacrylamide and N,N-dimethylacrylamide), vinyl formamide, vinyl pyrrolidone, alkoxylated (meth)acrylate, such as hydroxyethyl(meth)acrylate,
- 20 hydroxypropyl(meth)acrylate, and their higher ethoxylated or propoxylated versions such as behenyl polyethoxy methacrylate, e.g. sold as Sipomer BEM ex Rhodia, of the formula (V):



wherein R^{15} is hydrogen, or methyl and R^{16} is hydrogen, methyl or ethyl, and X is from 1 to 150;


5

- (ii) hydrophobic neutral monomers such as vinyl acetate and its higher homologs, alkyl(meth)acrylates (e.g. methyl methacrylate, butyl acrylate and ethyl acrylate), styrene and its derivatives, methyl vinyl ether, Sipomer WAM and WAM II from Rhodia, glycidyl methacrylate, and many commercial speciality monomers used in conventional latexes (e.g. AAEM, other from DGY). Of the above mentioned monomers, those which are hydrolysable in the wash medium are especially preferred (e.g. vinyl acetate and its derivatives)

15

- (iii) hydrophilic neutral with potentially cationic functional groups. The said functional groups have a pK lower than 12 preferably between 5 to 11 and most preferably between 6 to 10. The monomers above which comprise an amino group are especially preferred. The most preferred uncharged monomers are of formula (VI):

20



wherein R^{17} is hydrogen or C_1 to C_6 alkyl, Z^4 is O (ether linkage), CO_2 (ester), or CONH (amide), R^{18} and R^{19} are independently hydrogen, C_1 to C_{10} with OH

25

and/or NH_2 groups at the end or any position along this alkyl chain, and v is from 1 to 6.

Typical suitable such uncharged monomer units independently may be selected as one
5 or more of the optional uncharged monomer unit(s) denoted by formula (III).

Synthesis of the Polymer

10 The polymer used in the present invention is preferably obtained from respective monomers corresponding to anionic monomer units (a), cationic monomer units (b) and optionally, neutral (uncharged) monomer units (c), each respectively being ethylenically unsaturated. The different available means of copolymerising such ethylenically
15 unsaturated monomers will be well known to those skilled in the art of polymer chemistry. Depending on the order of addition of reactants, the resulting polymers may be block, random or mixed block/random copolymers.

Surfactants

20 Compositions according to the first aspect of the invention must also comprise one or more surfactants suitable for use in laundry wash and/or rinsing products. In the most general sense, these may be chosen from one or more of soap and non-soap anionic, cationic, nonionic, amphoteric and zwitterionic surface-active compounds and mixtures
25 thereof. Many suitable surface-active compounds are available and are fully described in the literature, for example, in "Surface-Active Agents and Detergents", Volumes I and II, by Schwartz, Perry and Berch.

For those compositions intended as laundry wash products, preferably, the surfactant(s)
30 is/are selected from one or more soaps and synthetic non-soap anionic and non-ionic compounds. Detergent compositions suitable for use in most automatic fabric washing

machines generally contain anionic non-soap surfactant, or non-ionic surfactant, or combinations of the two in any suitable ratio, optionally together with soap.

For example, laundry wash compositions of the invention may contain linear
5 alkylbenzene sulphonate anionic surfactants, particularly linear alkylbenzene sulphonates having an alkyl chain length of C₈-C₁₅. It is preferred if the level of linear alkylbenzene sulphonate is from 0 wt% to 30 wt%, more preferably 1 wt% to 25 wt%, most preferably from 2 wt% to 15 wt%.

10 The laundry wash compositions of the invention may additionally or alternatively contain one or more other anionic surfactants in total amounts corresponding to percentages quoted above for alkyl benzene sulphonates. Suitable anionic surfactants are well-known to those skilled in the art. These include primary and secondary alkyl
15 sulphates, particularly C₈-C₁₅ primary alkyl sulphates; alkyl ether sulphates; olefin sulphonates; alkyl xylene sulphonates; dialkyl sulphosuccinates; and fatty acid ester sulphonates. Sodium salts are generally preferred.

Some particular examples of such other anionic surfactants are:-

- 20 • alkyl ester sulphonates of the formula R-CH(SO₃M)-COOR', where R is a C₈-C₂₀, preferably C₁₀-C₁₆ alkyl radical, R' is a C₁-C₆, preferably C₁-C₃ alkyl radical, and M is an alkaline cation (sodium, potassium, lithium), substituted or non-substituted ammonium (methyl, dimethyl, trimethyl, tetramethyl ammonium, dimethyl piperidinium, etc.) or a derivative of an alkanol amine (monoethanol amine,
25 diethanol amine, triethanol amine, etc.);
- alkyl sulphates of the formula ROSO₃M, where R is a C₅-C₂₄, preferably C₁₀-C₁₈ alkyl or hydroxyalkyl radical, and M is a hydrogen atom or a cation as defined above, and their ethyleneoxy (EO) and/or propyleneoxy (PO) derivatives, having on average 0.5 to 30, preferably 0.5 to 10 EO and/or PO units;
- 30 • alkyl amide sulphates of the formula RCONHR'OSO₃M, where R is a C₂-C₂₂, preferably C₆-C₂₀ alkyl radical, R' is a C₂-C₃ alkyl radical, and M is a hydrogen atom

or a cation as defined above, and their ethyleneoxy (EO) and/or propyleneoxy (PO) derivatives, having on average 0.5 to 60 EO and/or PO units;

- the salts of C_8 - C_{24} , preferably C_{14} - C_{20} saturated or unsaturated fatty acids, C_8 - C_{22} primary or secondary alkyl sulphonates, alkyl glycerol sulphonates, the sulphonated polycarboxylic acids described in GB-A-1 082 179, paraffin sulphonates, N-acyl, N'-alkyl taurates, alkyl phosphates, isethionates, alkyl succinamates, alkyl sulphosuccinates, monoesters or diesters of sulphosuccinates, N-acyl sarcosinates, alkyl glycoside sulphates, polyethoxycarboxylates, the cation being an alkali metal (sodium, potassium, lithium), a substituted or non-substituted ammonium residue (methyl, dimethyl, trimethyl, tetramethyl ammonium, dimethyl piperidinium, etc.) or a derivative of an alkanol amine (monoethanol amine, diethanol amine, triethanol amine, etc.);
- sophorolipids, such as those in acid or lactone form, derived from 17-hydroxyoctadecenic acid;

The laundry wash compositions of the invention may contain non-ionic surfactant.

Nonionic surfactants that may be used include the primary and secondary alcohol ethoxylates, especially the C_8 - C_{20} aliphatic alcohols ethoxylated with an average of from 1 to 20 moles of ethylene oxide per mole of alcohol, and more especially the C_{10} - C_{15} primary and secondary aliphatic alcohols ethoxylated with an average of from 1 to 10 moles of ethylene oxide per mole of alcohol. Non-ethoxylated nonionic surfactants include alkylpolyglycosides, glycerol monoethers, and polyhydroxyamides (glucamide).

Some particular examples of such nonionic surfactants are:-

- polyalkoxylenated alkyl phenols (i.e. polyethyleneoxy, polypropyleneoxy, polybutyleneoxy), the alkyl substituent of which has from 6 to 12 C atoms and contains from 5 to 25 alkoxylenated units; examples are TRITON X-45, X-114, X-100 and X-102 marketed by Rohm & Haas Co., IGEPAL NP2 to NP17 made by Rhodia;

- C₈-C₂₂ polyalkoxylenated aliphatic alcohols containing 1 to 25 alkoxylenated (ethyleneoxy, propyleneoxy) units; examples are TERGITOL 15-S-9, TERGITOL 24-L-6 NMW marketed by Union Carbide Corp., NEODOL 45-9, NEODOL 23-65, NEODOL 45-7, NEODOL 45-4 marketed by Shell Chemical Co., KYRO EOB
5 marketed by The Procter & Gamble Co., SYNPERONIC A3 to A9 made by ICI, RHODASURF IT, DB and B made by Rhodia;
- the products resulting from the condensation of ethylene oxide or propylene oxide with propylene glycol, ethylene glycol, with a molecular weight in the order of 2000 to 10,000, such as the PLURONIC products marketed by BASF;
- 10 • the products resulting from the condensation of ethylene oxide or propylene oxide with ethylene diamine, such as the TETRONIC products marketed by BASF;
- C₈-C₁₈ ethoxyl and/or propoxyl fatty acids containing 5 to 25 ethyleneoxy and/or propyleneoxy units;
- C₈-C₂₀ fatty acid amides containing 5 to 30 ethyleneoxy units;
- 15 • ethoxylated amines containing 5 to 30 ethyleneoxy units;
- alkoxyated amidoamines containing 1 to 50, preferably 1 to 25 and in particular 2 to 20 alkyleneoxy (preferably ethyleneoxy) units;
- amine oxides such as the oxides of alkyl C₁₀-C₁₈ dimethylamines, the oxides of alkoxy C₈-C₂₂ ethyl dihydroxy ethylamines;
- 20 • alkoxyated terpene hydrocarbons such as ethoxylated and/or propoxylated α- or β-pinenes, containing 1 to 30 ethyleneoxy and/or propyleneoxy units;
- alkylpolyglycosides obtainable by condensation (for example by acid catalysis) of glucose with primary fatty alcohols (e.g. US-A-3 598 865; US-A-4 565 647; EP-A-132 043; EP-A-132 046) having a C₄-C₂₀, preferably C₈-C₁₈ alkyl group and an
25 average number of glucose units in the order of 0.5 to 3, preferably in the order of 1.1 to 1.8 per mole of alkylpolyglycoside (APG), particularly those having
 - * a C₈-C₁₄ alkyl group and on average 1.4 glucose units per mole
 - * a C₁₂-C₁₄ alkyl group and on average 1.4 glucose units per mole
 - * a C₈-C₁₄ alkyl group and on average 1.5 glucose units per mole
 - 30 * a C₈-C₁₀ alkyl group and on average 1.6 glucose units per mole

marketed under the names GLUCOPON 600 EC[®], GLUCOPON 600 CSUP[®], GLUCOPON 650 EC[®] and GLUCOPON 225 CSUP[®] respectively and made by HENKEL;

- 5 It is preferred if the level of total non-ionic surfactant is from 0 wt% to 30 wt%, preferably from 1 wt% to 25 wt%, most preferably from 2 wt% to 15 wt%.

Another class of suitable surfactants comprises certain mono-long chain-alkyl cationic surfactants for use in main-wash laundry compositions according to the invention.

- 10 Cationic surfactants of this type include quaternary ammonium salts of the general formula $R_1R_2R_3R_4N^+ X^-$ wherein the R groups are long or short hydrocarbon chains, typically alkyl, hydroxyalkyl or ethoxylated alkyl groups, and X is a counter-ion (for example, compounds in which R₁ is a C₈-C₂₂ alkyl group, preferably a C₈-C₁₀ or C₁₂-C₁₄ alkyl group, R₂ is a methyl group, and R₃ and R₄, which may be the same or different,
15 are methyl or hydroxyethyl groups); and cationic esters (for example, choline esters).

- The choice of surface-active compound (surfactant), and the amount present in the laundry wash compositions according to the invention, will depend on the intended use of the detergent composition. In fabric washing compositions, different surfactant
20 systems may be chosen, as is well known to the skilled formulator, for handwashing products and for products intended for use in different types of washing machine. The total amount of surfactant present will also depend on the intended end use and may be as high as 60 wt%, for example, in a composition for washing fabrics by hand. In compositions for machine washing of fabrics, an amount of from 5 to 40 wt% is
25 generally appropriate. Typically the compositions will comprise at least 2 wt% surfactant e.g. 2-60%, preferably 15-40% most preferably 25-35%.

- In the case of laundry rinse compositions according to the invention the surfactant(s) is/are preferably selected from fabric conditioning agents. In fact, conventional fabric
30 conditioning agent may be used. These conditioning agents may be cationic or non-ionic. If the fabric conditioning compound is to be employed in a main wash detergent

composition the compound will typically be non-ionic. If used in the rinse phase, they will typically be cationic. They may for example be used in amounts from 0.5% to 35%, preferably from 1% to 30% more preferably from 3% to 25% by weight of the composition.

5

Preferably the fabric conditioning agent(s) have two long chain alkyl or alkenyl chains each having an average chain length greater than or equal to C₁₆. Most preferably at least 50% of the long chain alkyl or alkenyl groups have a chain length of C₁₈ or above. It is preferred if the long chain alkyl or alkenyl groups of the fabric conditioning agents

10

The fabric conditioning agents are preferably compounds that provide excellent softening, and are characterised by a chain melting L β to L α transition temperature greater than 25°C, preferably greater than 35°C, most preferably greater than 45°C. This

15

L β to L α transition can be measured by DSC as defined in "Handbook of Lipid Bilayers, D Marsh, CRC Press, Boca Raton, Florida, 1990 (pages 137 and 337).

Substantially insoluble fabric conditioning compounds in the context of this invention are defined as fabric conditioning compounds having a solubility less than 1×10^{-3} wt %

20

in demineralised water at 20°C. Preferably the fabric softening compounds have a solubility less than 1×10^{-4} wt %, most preferably less than 1×10^{-8} to 1×10^{-6} .

Preferred cationic fabric softening agents comprise a substantially water insoluble quaternary ammonium material comprising a single alkyl or alkenyl long chain having an average chain length greater than or equal to C₂₀ or, more preferably, a compound

25

comprising a polar head group and two alkyl or alkenyl chains having an average chain length greater than or equal to C₁₄.

Preferably, the cationic fabric softening agent is a quaternary ammonium material or a quaternary ammonium material containing at least one ester group. The quaternary

30

ammonium compounds containing at least one ester group are referred to herein as ester-linked quaternary ammonium compounds.

As used in the context of the quaternary ammonium cationic fabric softening agents, the term 'ester group', includes an ester group which is a linking group in the molecule.

- 5 It is preferred for the ester-linked quaternary ammonium compounds to contain two or more ester groups. In both monoester and the diester quaternary ammonium compounds it is preferred if the ester group(s) is a linking group between the nitrogen atom and an alkyl group. The ester groups(s) are preferably attached to the nitrogen atom via another hydrocarbyl group.

10

- Also preferred are quaternary ammonium compounds containing at least one ester group, preferably two, wherein at least one higher molecular weight group containing at least one ester group and two or three lower molecular weight groups are linked to a common nitrogen atom to produce a cation and wherein the electrically balancing anion is a halide, acetate or lower alkylsulphate ion, such as chloride or methosulphate. The higher molecular weight substituent on the nitrogen is preferably a higher alkyl group, containing 12 to 28, preferably 12 to 22, e.g. 12 to 20 carbon atoms, such as coco-alkyl, tallowalkyl, hydrogenated tallowalkyl or substituted higher alkyl, and the lower molecular weight substituents are preferably lower alkyl of 1 to 4 carbon atoms, such as methyl or ethyl, or substituted lower alkyl. One or more of the said lower molecular weight substituents may include an aryl moiety or may be replaced by an aryl, such as benzyl, phenyl or other suitable substituents.

20

- Preferably the quaternary ammonium material is a compound having two C_{12} - C_{22} alkyl or alkenyl groups connected to a quaternary ammonium head group via at least one ester link, preferably two ester links or a compound comprising a single long chain with an average chain length equal to or greater than C_{20} .


25

- More preferably, the quaternary ammonium material comprises a compound having two long chain alkyl or alkenyl chains with an average chain length equal to or greater than C_{14} . Even more preferably each chain has an average chain length equal to or greater

30

than C₁₆. Most preferably at least 50% of each long chain alkyl or alkenyl group has a chain length of C₁₈. It is preferred if the long chain alkyl or alkenyl groups are predominantly linear.

- 5 The most preferred type of ester-linked quaternary ammonium material that can be used in laundry rinse compositions according to the invention is represented by the formula (A):



wherein T is $\begin{array}{c} O \\ || \end{array}$ or $\begin{array}{c} O \\ || \end{array}$; each R²⁰ group is independently selected from C₁₋₄ alkyl, hydroxyalkyl or C₂₋₄ alkenyl groups; and wherein each R²¹ group is independently selected from C₈₋₂₈ alkyl or alkenyl groups; Y⁻ is any suitable counter-ion, i.e. a halide, acetate or lower alkylsulphate ion, such as chloride or methosulphate;

20

w is an integer from 1-5 or is 0; and

25 y is an integer from 1-5.

It is especially preferred that each R²⁰ group is methyl and w is 1 or 2.


It is advantageous for environmental reasons if the quaternary ammonium material is

30 biologically degradable.

Preferred materials of this class such as 1,2 bis[hardened tallowyloxy]-3-trimethylammonium propane chloride and their method of preparation are, for example, described in US-A-4 137 180. Preferably these materials comprise small amounts of the corresponding monoester as described in US-A-4 137 180 for example 1-hardened
 5 tallowyloxy-2-hydroxy-3-trimethylammonium propane chloride.

Another class of preferred ester-linked quaternary ammonium materials for use in laundry rinse compositions according to the invention can be represented by the formula:

10




20 wherein T is $\begin{array}{c} O \\ || \end{array}$ or $\begin{array}{c} O \\ || \end{array}$ -O-C- or -C-O-; and

wherein R^{20} , R^{21} , w, and Y^- are as defined above.

Of the compounds of formula (B), di-(tallowyloxyethyl)-dimethyl ammonium chloride, available from Hoechst, is the most preferred. Di-(hardened tallowyloxyethyl)dimethyl ammonium chloride, ex Hoechst and di-(tallowyloxyethyl)-methyl hydroxyethyl methosulphate are also preferred.

30

Another preferred class of quaternary ammonium cationic fabric softening agent is defined by formula (C):-



5

where R^{20} , R^{21} and Y^{-} are as hereinbefore defined.

A preferred material of formula (C) is di-hardened tallow-diethyl ammonium chloride, sold under the Trademark Arquad 2HT.

10

The optionally ester-linked quaternary ammonium material may contain optional additional components, as known in the art, in particular, low molecular weight solvents, for instance isopropanol and/or ethanol, and co-actives such as nonionic softeners, for example fatty acid or sorbitan esters.

15

Detergency Builders

The compositions of the invention, when used as laundry wash compositions, will generally also contain one or more detergency builders. The total amount of detergency builder in the compositions will typically range from 5 to 80 wt%, preferably from 10 to 60 wt%.

Inorganic builders that may be present include sodium carbonate, if desired in combination with a crystallisation seed for calcium carbonate, as disclosed in GB 1 437 950 (Unilever); crystalline and amorphous aluminosilicates, for example, zeolites as disclosed in GB 1 473 201 (Henkel), amorphous aluminosilicates as disclosed in GB 1 473 202 (Henkel) and mixed crystalline/amorphous aluminosilicates as disclosed in

25

GB 1 470 250 (Procter & Gamble); and layered silicates as disclosed in EP 164 514B (Hoechst). Inorganic phosphate builders, for example, sodium orthophosphate, pyrophosphate and tripolyphosphate are also suitable for use with this invention.

- 5 The compositions of the invention preferably contain an alkali metal, preferably sodium, aluminosilicate builder. Sodium aluminosilicates may generally be incorporated in amounts of from 10 to 70% by weight (anhydrous basis), preferably from 25 to 50 wt%.
- 10 The alkali metal aluminosilicate may be either crystalline or amorphous or mixtures thereof, having the general formula: $0.8-1.5 \text{ Na}_2\text{O} \cdot \text{Al}_2\text{O}_3 \cdot 0.8-6 \text{ SiO}_2$.

- These materials contain some bound water and are required to have a calcium ion exchange capacity of at least 50 mg CaO/g. The preferred sodium aluminosilicates
- 15 contain 1.5-3.5 SiO_2 units (in the formula above). Both the amorphous and the crystalline materials can be prepared readily by reaction between sodium silicate and sodium aluminate, as amply described in the literature. Suitable crystalline sodium aluminosilicate ion-exchange detergency builders are described, for example, in GB 1 429 143 (Procter & Gamble). The preferred sodium aluminosilicates of this type are the
- 20 well-known commercially available zeolites A and X, and mixtures thereof.

- The zeolite may be the commercially available zeolite 4A now widely used in laundry detergent powders. However, according to a preferred embodiment of the invention, the zeolite builder incorporated in the compositions of the invention is maximum aluminium
- 25 zeolite P (zeolite MAP) as described and claimed in EP 384 070A (Unilever). Zeolite MAP is defined as an alkali metal aluminosilicate of the zeolite P type having a silicon to aluminium ratio not exceeding 1.33, preferably within the range of from 0.90 to 1.33, and more preferably within the range of from 0.90 to 1.20.

Especially preferred is zeolite MAP having a silicon to aluminium ratio not exceeding 1.07, more preferably about 1.00. The calcium binding capacity of zeolite MAP is generally at least 150 mg CaO per g of anhydrous material.

- 5 Organic builders that may be present include polycarboxylate polymers such as polyacrylates, acrylic/maleic copolymers, and acrylic phosphinates; monomeric polycarboxylates such as citrates, gluconates, oxydisuccinates, glycerol mono-, di and trisuccinates, carboxymethyloxy succinates, carboxymethyloxymalonates, dipicolinates, hydroxyethyliminodiacetates, alkyl- and alkenylmalonates and succinates; and
10 sulphonated fatty acid salts. This list is not intended to be exhaustive.

- Especially preferred organic builders are citrates, suitably used in amounts of from 5 to 30 wt%, preferably from 10 to 25 wt%; and acrylic polymers, more especially acrylic/maleic copolymers, suitably used in amounts of from 0.5 to 15 wt%, preferably
15 from 1 to 10 wt%.

Builders, both inorganic and organic, are preferably present in alkali metal salt, especially sodium salt, form.

20

Bleaches

- Laundry wash compositions according to the invention may also suitably contain a bleach system. Fabric washing compositions may desirably contain peroxy bleach
25 compounds, for example, inorganic persalts or organic peroxyacids, capable of yielding hydrogen peroxide in aqueous solution.

- Suitable peroxy bleach compounds include organic peroxides such as urea peroxide, and inorganic persalts such as the alkali metal perborates, percarbonates, perphosphates, persilicates and persulphates. Preferred inorganic persalts are sodium perborate
30 monohydrate and tetrahydrate, and sodium percarbonate.

Especially preferred is sodium percarbonate having a protective coating against destabilisation by moisture. Sodium percarbonate having a protective coating comprising sodium metaborate and sodium silicate is disclosed in GB 2 123 044B (Kao).

5

The peroxy bleach compound is suitably present in an amount of from 0.1 to 35 wt%, preferably from 0.5 to 25 wt%. The peroxy bleach compound may be used in conjunction with a bleach activator (bleach precursor) to improve bleaching action at low wash temperatures. The bleach precursor is suitably present in an amount of from 0.1 to 8 wt%, preferably from 0.5 to 5 wt%.

10

Preferred bleach precursors are peroxycarboxylic acid precursors, more especially peracetic acid precursors and pernonanoic acid precursors. Especially preferred bleach precursors suitable for use in the present invention are N,N,N',N'-tetracetyl

15

ethylenediamine (TAED) and sodium nonanoyloxybenzene sulphonate (SNOBS). The novel quaternary ammonium and phosphonium bleach precursors disclosed in US 4 751 015 and US 4 818 426 (Lever Brothers Company) and EP 402 971A (Unilever), and the cationic bleach precursors disclosed in EP 284 292A and EP 303 520A (Kao) are also of interest.

20

The bleach system can be either supplemented with or replaced by a peroxyacid. examples of such peracids can be found in US 4 686 063 and US 5 397 501 (Unilever). A preferred example is the imido peroxycarboxylic class of peracids described in EP A 325 288, EP A 349 940, DE 382 3172 and EP 325 289. A particularly preferred example is phthalimido peroxy caproic acid (PAP). Such peracids are suitably present at 0.1 - 12%, preferably 0.5 - 10%.

25

A bleach stabiliser (transition metal sequestrant) may also be present. Suitable bleach stabilisers include ethylenediamine tetra-acetate (EDTA), the polyphosphonates such as Dequest (Trade Mark) and non-phosphate stabilisers such as EDDS (ethylene diamine

30

di-succinic acid). These bleach stabilisers are also useful for stain removal especially in products containing low levels of bleaching species or no bleaching species.

5 An especially preferred bleach system comprises a peroxy bleach compound (preferably sodium percarbonate optionally together with a bleach activator), and a transition metal bleach catalyst as described and claimed in EP 458 397A, EP 458 398A and EP 509 787A (Unilever).

10 Enzymes

Laundry wash compositions according to the invention may also contain one or more enzyme(s). Suitable enzymes include the proteases, amylases, cellulases, oxidases, peroxidases and lipases usable for incorporation in detergent compositions. Preferred
15 proteolytic enzymes (proteases) are catalytically active protein materials which degrade or alter protein types of stains when present as in fabric stains in a hydrolysis reaction. They may be of any suitable origin, such as vegetable, animal, bacterial or yeast origin.

Proteolytic enzymes or proteases of various qualities and origins and having activity in
20 various pH ranges of from 4-12 are available and can be used in the instant invention. Examples of suitable proteolytic enzymes are the subtilisins which are obtained from particular strains of B. Subtilis B. licheniformis, such as the commercially available subtilisins Maxatase (Trade Mark), as supplied by Gist Brocades N.V., Delft, Holland, and Alcalase (Trade Mark), as supplied by Novo Industri A/S, Copenhagen, Denmark.

25 Particularly suitable is a protease obtained from a strain of Bacillus having maximum activity throughout the pH range of 8-12, being commercially available, e.g. from Novo Industri A/S under the registered trade-names Esperase (Trade Mark) and Savinase (Trade-Mark). The preparation of these and analogous enzymes is described in GB 1 243
30 785. Other commercial proteases are Kazusase (Trade Mark obtainable from

Showa-Denko of Japan), Optimase (Trade Mark from Miles Kali-Chemie, Hannover, West Germany), and Superase (Trade Mark obtainable from Pfizer of U.S.A.).

Detergency enzymes are commonly employed in granular form in amounts of from about
5 0.1 to about 3.0 wt%. However, any suitable physical form of enzyme may be used.

Other Optional Ingredients

10 The compositions of the invention may contain alkali metal, preferably sodium carbonate, in order to increase detergency and ease processing. Sodium carbonate may suitably be present in amounts ranging from 1 to 60 wt%, preferably from 2 to 40 wt%. However, compositions containing little or no sodium carbonate are also within the scope of the invention.

15 Powder flow may be improved by the incorporation of a small amount of a powder structurant, for example, a fatty acid (or fatty acid soap), a sugar, an acrylate or acrylate/maleate copolymer, or sodium silicate. One preferred powder structurant is fatty acid soap, suitably present in an amount of from 1 to 5 wt%.

20 Yet other materials that may be present in detergent compositions of the invention include sodium silicate; antiredeposition agents such as cellulosic polymers; inorganic salts such as sodium sulphate; lather control agents or lather boosters as appropriate; proteolytic and lipolytic enzymes; dyes; coloured speckles; perfumes; foam controllers;
25 fluorescers and decoupling polymers. This list is not intended to be exhaustive.

It is often advantageous if soil release or soil suspending polymers are present, for example in amounts in the order of 0.01% to 10%, preferably in the order of 0.1% to 5% and in particular in the order of 0.2% to 3% by weight, such as
30 - cellulose derivatives such as cellulose hydroxyethers, methyl cellulose, ethyl cellulose, hydroxypropyl methyl cellulose, hydroxybutyl methyl cellulose;

- polyvinyl esters grafted onto polyalkylene backbones, such as polyvinyl acetates grafted onto polyoxyethylene backbones (EP-A-219 048);
- polyvinyl alcohols;
- polyester copolymers based on ethylene terephthalate and/or propylene terephthalate units and polyethyleneoxy terephthalate units, with a molar ratio (number of units) of ethylene terephthalate and/or propylene terephthalate / (number of units) polyethyleneoxy terephthalate in the order of 1/10 to 10/1, the polyethyleneoxy terephthalate units having polyethyleneoxy units with a molecular weight in the order of 300 to 10,000, with a molecular weight of the copolyester in the order of 1000 to 100,000;
- polyester copolymers based on ethylene terephthalate and/or propylene terephthalate units and polyethyleneoxy and/or polypropyleneoxy units, with a molar ratio (number of units) of ethylene terephthalate and/or propylene terephthalate / (number of units) polyethyleneoxy and/or polypropyleneoxy in the order of 1/10 to 10/1, the polyethyleneoxy and/or polypropyleneoxy units having a molecular weight in the order of 250 to 10,000, with a molecular weight of the copolyester in the order of 1000 to 100,000 (US-A-3 959 230, US-A-3 962 152, US-A-3 893 929, US-A-4 116 896, US-A-4 702 857, US-A-4 770 666, EP-A-253 567, EP-A-201 124);
- copolymers of ethylene or propylene terephthalate / polyethyleneoxy terephthalate comprising sulphoisophthaloyl units in their chain (US-A-4 711 730, US-A-4 702 857, US-A-4 713 194);
- terephthalic copolyester oligomers having polyalkyleneoxyalkyl sulphonate/sulphoaroyl terminal groups and optionally containing sulphoisophthaloyl units in their chain (US-A-4 721 580, US-A-5 415 807, US-A-4 877 896, US-A-5 182 043, US-A-5 599 782, US-A-4 764 289, EP-A-311 342, WO92/04433, WO97/42293);
- sulphonated terephthalic copolyesters with a molecular weight less than 20,000, obtained e.g. from a diester of terephthalic acid, isophthalic acid, a diester of sulphoisophthalic acid and a diol, in particular ethylene glycol (WO95/32997);
- polyurethane polyesters, obtained by reaction of a polyester with a molecular weight of 300 to 4000, obtained from a terephthalic acid diester, possibly a

- 5 sulphisophthalic acid diester and a diol, on a prepolymer with isocyanate terminal groups, obtained from a polyethyleneoxy glycol with a molecular weight of 600 to 4000 and a diisocyanate (US-A-4 201 824);
- 5 sulphonated polyester oligomers obtained by sulphonation of an oligomer derived from ethoxylated allyl alcohol, dimethyl terephthalate and 1,2-propylene diol, having 1 to 4 sulphonate groups (US-A-4 968 451).

Use

10

The composition when diluted in the wash liquor (during a typical wash cycle) will typically give a pH of the wash liquor from 7 to 11, preferably from 7 to 10.5, for a wash product. Treatment of a fabric with a soil-release polymer in accordance with the second aspect of the present invention can be made by any suitable method such as washing, soaking or rinsing.

15

Typically the treatment will involve a washing or rinsing method such as treatment in the main wash or rinse cycle of a washing machine and involves contacting the fabric with an aqueous medium comprising the composition according to the first aspect of the present invention.

20

Product Form

25 Compositions according to the first aspect of the present invention may be formulated in any convenient form, for example as powders, liquids (aqueous or non-aqueous) or tablets.

30 Particulate detergent compositions are suitably prepared by spray-drying a slurry of compatible heat-insensitive ingredients, and then spraying on or post-dosing those ingredients unsuitable for processing via the slurry. The skilled detergent formulator

will have no difficulty in deciding which ingredients should be included in the slurry and which should not.

Particulate detergent compositions of the invention preferably have a bulk density of at least 400 g/l, more preferably at least 500 g/l. Especially preferred compositions have bulk densities of at least 650 g/litre, more preferably at least 700 g/litre.

Such powders may be prepared either by post-tower densification of spray-dried powder, or by wholly non-tower methods such as dry mixing and granulation; in both cases a high-speed mixer/granulator may advantageously be used. Processes using high-speed mixer/granulators are disclosed, for example, in EP 340 013A, EP 367 339A, EP 390 251A and EP 420 317A (Unilever).

Liquid detergent compositions can be prepared by admixing the essential and optional ingredients thereof in any desired order to provide compositions containing components in the requisite concentrations. Liquid compositions according to the present invention can also be in compact form which means it will contain a lower level of water compared to a conventional liquid detergent.

The present invention will now be explained in more detail by way of the following non-limiting examples.

Examples

25

The following evaluation experiments were carried-out using base formulations of detergent powders as follows (hereinafter called "Composition A" and "Composition B").

30

Composition A

		wt %
	LAS ¹	24.0
5	STPP ²	14.5
	Sodium silicate	7.0
	Sodium sulphate	17.7
	Sodium perborate	8.0
	TAED ³	2.4
10	Enzymes and minors	balance

Composition B

		wt %
	LAS ¹	23.00
15	STPP ²	14.50
	Sodium carbonate	17.50
	Sodium sulphate	28.52
	Sodium silicate	7.00
20	Enzymes and minors	balance

1. C₁₂ - alkylbenzene sulphonate, sodium salt
2. Sodium tripolyphosphate
3. Tetra acetyl ethylene diamine, bleach activator

25

Experimental Procedures and MaterialsExample 1Evaluation Procedure

- 30 1. Pretreatment:
 - with the polymer (when present) at 200mg/l


- in a washing solution at 2.1g/l Composition A (Brazilian detergent powder), 5°TH
 - 23°C, 20 minutes, 100rpm, tergotometer
 - rinse in 1 litre water 5 minutes
 - drying on the drying drum (drum temperature is 95°C; the swatches make ~3 turns
- 5 over ~2 minutes; temperature of fabric taken off drum ~35°C)
2. Stain with 4 drops of dirty motor oil (DMO) diluted to 15% in toluene; dry at 40°C for 1 hour
- 10 3. Wash with 2.1g/l Composition A, 5°TH, 20 minutes at 23°C, rinse 5 minutes in 1 litre tap water, dry on the drying drum

Polymers


15

Two series of polyampholytes based on monomers of acrylic acid (AA) and quaternary methacrylamides MAPTAC or diQUAT were studied. The polymer characteristics are given in Tables 1-3, in which the following abbreviations are used for the constituent monomers:-


DIQUAT




AA



MAPTAC



DMAEMA






Table 1: AA/MAPTAC polyampholytes

Polymer Example	AA mole	MAPTAC mole	active /%	viscosity /cps
1	50	50	20.6 %	36 000
2	70	30	20.1 %	46 500
3	90	10	15.2 %	36 000
4	95	5	15.1 %	25 000

5

Table 2: AA/DIQUAT polyampholytes

Polymer Example	AA mole	DIQUAT mole	active /%	viscosity /cps
5	80	20	17.1 %	37 250
6	67	33	20.3 %	6 150
7	50	50	20.5 %	840

Table 3: Ter-polymer DMAEMA/MA/MES

Polymer Example	DMAEMA mole	MA mole	MES mole	molecular weight
8	40	30	30	low
9	40	40	20	
10	40	50	10	
11	40	30	30	high
12	40	40	20	
13	40	50	10	

5 Results

In the following tables, the polymer was added in the amount specified to the wash solution based on Formulation A, as specified above

10 Table 4: AA/MAPTAC Polymers

Polymer Example	Amount mg/l	ΔR
---	---	13.5
1	200	15.6
2	200	17.5
3	200	15.3
4	200	14.9

Table 4: AA/DIQUAT Polymers

Polymer Example	Amount mg/l	ΔR	Amount mg/l	ΔR
---	---	13.1	---	13.1
5	100	17.7	200	17.5
6	100	9.5	200	8.2
7	---	---	200	7.1

5 Table 5: Example 5 - Effect of Dosage

Amount of Example 5 Polymer (mg/l)	ΔR
0	13.0
20	16.2
40	16.4
60	18.4
80	17.9
100	18.3

Table 6: DMAEMA/MAA/MES

Polymer Example	Amount mg/l	ΔR	Amount mg/l	ΔR
---	---	13.4	---	13.4
11	100	15.3	200	18.7
12	100	16.6	200	19.5
13	---	---	200	14.7

Example 2Evaluation procedure







Conditions: tergotometer, 100rpm, 23°C.

- 5 1. PRE-WASH: 6 3"x3" desized cotton squares, in 1 litre of wash liquor
(liquor: cloth ca. 200:1)
- 10 wash liquor: - 2.1g of detergent (Composition B)
- 200ppm of polymer
- water: 1 litre water 12°FH
- agitated for 20 mins
- 15 wash liquor decanted off
- Rinse: 1 litre of 12°FH water (as before)
- Agitated for 5 mins
- 20 2. STAINING: dirty motor oil (DMO) diluted to 15 wt. %
- 25 4 drops of diluted DMO applied from a burette to each 3"x3" square
Left to dry on racks in an oven (40°C) for 1 hour
3. MAIN WASH & rinse: as pre-wash except polymer is at 50ppm.
- 30 4. ANALYSIS: results are obtained by extracting R460 values (reflectance values at 460 nm) of the cloths
- 35 1. before staining (R_{clean})
2. after staining (R_{stain})
3. after final washing (R_{washed})
- delta (Δ) R is calculated for all samples including control (no polymer treatment):
 $R_{\text{washed}} - R_{\text{stain}}$
- 40 $\Delta\Delta R$ is then calculated for quick comparison to the control
- $\Delta R_{\text{polymer}} - \Delta R_{\text{control}}$
- 45


Polymers

- 5 Various polyampholytes were prepared and studied. The polymer compositions and test results are given in Table 7, in which the following abbreviations are used for the constituent monomers:-

NEUTRAL MONOMERS

Name	Structure
N-Vinylformamide (NVF)	
t-Butylaminoethyl methacrylate (t-BuAEMA)	
Dimethylaminopropyl-methacrylamide (DMAPMA)	
Diethylaminoethyl Methacrylate (DEAEMA)	
Dimethylaminoethyl Acrylate (DMAEA)	
N, N-dimethylacrylamide (DMA)	$(\text{CH}_3)_2\text{NC}(\text{O})\text{CH}=\text{CH}_2$
Vinyl acetate (VAc)	$\text{CH}_2=\text{CHOC}(\text{O})\text{CH}_3$
Hydroxylpropyl acrylate (HPA)	$\text{HOCH}_2\text{CH}_2\text{CH}_2\text{OC}(\text{O})\text{CH}=\text{CH}_2$
Hydroxyethyl acrylate (HEA)	$\text{HOCH}_2\text{CH}_2\text{OC}(\text{O})\text{CH}=\text{CH}_2$
Hydroxyethyl methacrylate (HEMA)	$\text{HOCH}_2\text{CH}_2\text{OC}(\text{O})\text{C}(\text{CH}_3)=\text{CH}_2$
Ethylhexylacrylate (EHA)	

ANIONIC MONOMERS

Name	Structure
Acrylic Acid (AA)	$\text{CH}_2=\text{CHCO}_2\text{H}$
Methacrylic Acid (MAA)	$\text{CH}_2=\text{C}(\text{CH}_3)\text{CO}_2\text{H}$
Acrylamidomethylpropanesulfonic acid (AMPS)	

5 CATIONIC MONOMERS.



Name	Structure
Methacrylamidopropyl-trimethylammonium chloride/methylsulfate MAPTAC/MES	
Diquat	

Table 7

neutral (0)	anionic (-ve)	cationic (+ve)	% neutral	% anionic	% cationic	$\Delta\Delta R$ DMO
VP	AMPS	Diquat	40	40	20	0.2
V-Ac	AMPS	MAPTAC	40	40	20	1.2
t-BuAEMA	AMPS	MAPTAC	40	40	20	0.9
PEG2000-MA	MAA	MAPTAC	30	46.7	23.3	1.9
NVF	AMPS	MAPTAC	40	40	20	1.1
NVF	AMPS	Diquat	44	44	11	1.6
HPA	AMPS	MAPTAC	40	40	20	2.8
HPA	AA	MAPTAC	40	40	20	2.6
HPA	AMPS	Diquat	44	44	11	4.1
HPA	AA	Diquat	44	44	11	0.2
HEMA-10	MAA	MES	40	40	20	3.5
DMAEMA						
HEA	AMPS	Diquat	44	44	11	1.4
HEA	AA	DADMAC	40	40	20	3.0
EHA	AA	MES	24	61	15	0.1
DMAPMA	AMPS	MAPTAC	40	40	20	1.3
DMAPMA	AA	MAPTAC	20	60	20	0.4
DMAPMA	MAA	MAPTAC	20	60	20	0.1
DMAPMA	AMPS	Diquat	44	44	11	3.5
DMAEMA	MAA	MES	40	50	10	4.5
DMAEMA	MAA	MES	40	40	20	1.3
DMAEMA	AMPS	MES	40	30	30	0.9
DMAEMA	AMPS	MES	40	50	10	0.4
DMAEMA	MAA	MAPTAC	40	50	10	2.7
DMAEMA	MAA	MAPTAC	20	66.6	13.3	1.4
DMAEMA	MAA	MAPTAC	10	75	15	1.5
DMAEMA	MAA	DADMAC	20	73.3	6.7	1.5
DMA	MAA	MAPTAC	40	40	20	4.2
DMA	AMPS	MAPTAC	40	40	20	3.6
DMA	AMPS	Diquat	44	44	11	4.9
DEAEMA	AMPS	MAPTAC	40	40	20	0.7
DEAEMA	AMPS	Diquat	44	44	11	3.4
-	MAA	DADMAC	0	91.6	8.3	2.6

Example 3Washing-machine tests5 Evaluation procedure

- Conditions: Brazilian "Brastemp" washing-machine
 81 mins wash-time
 41 litres intake volume
 6°FH water
 25°C
 2.1g/l detergent formulation (composition B)
 2kg load weight
- 15 Pre-wash : 4 pre-washes at 50ppm polymer concentration
 Main-wash : single wash at 50ppm polymer concentration
- Staining
- 20 5 stains used,
 1. dirty motor oil (DMO), neat oil pipetted onto fabric
 2. artificial sebum dyed with Macrolex B dye (obtained from Bayer), painted onto fabric whilst hot*.
 3. clay (50% wt suspension in water), brushed onto fabric
 4. wine, pipetted neat onto fabric
 5. tea, pipetted neat onto fabric
- 25 * The artificial sebum was prepared by adding the following ingredients in the ratios given, heating to 60° C in a water bath and mixing until all the ingredients are combined:-
- 30 Sebum:
 Fatty Acids (Prifac 9484) 23.34%
 Tripalmitin (C16:0) 23.34%
 Brominated Olive Oil 23.34%
 Squalene 0.58%
 Cholesterol 4.92%
 Ceto Stearyl Stearate (Estol 1481) 14.42%
- 35
- 40 Once all the sebum was liquid, 0.08g of Macrolex B dye was added to 100g of sebum and the mixture was stirred whilst hot until the dye had mixed thoroughly.

Results**On cotton**

							$\Delta\Delta R$ DMO	$\Delta\Delta R$ sebum	$\Delta\Delta R$ clay	$\Delta\Delta R$ wine	$\Delta\Delta R$ tea
	DMAEMA	MAA	MES	40	40	20	0.1	3.5	2.2	1.3	0.4
	DMAEMA	MAA	MES	40	50	10	-0.2	-0.8	-0.4	0.8	0.2
	HEMA-10; MAEMA	MAA	MES	10; 30	40	20	0.0	1.3	-1.8	1.0	0.4
	DMAEMA	MAA	MAPTAC	20	66.6	13.3	2.1	0.7	-0.6	1.2	1.7
	DMAEMA	MAA	MAPTAC	40	50	10	1.8	2.7	0.8	0.9	2.1

5

On polyester-cotton

							$\Delta\Delta R$ DMO	$\Delta\Delta R$ sebum	$\Delta\Delta R$ clay	$\Delta\Delta R$ wine	$\Delta\Delta R$ tea
	DMAEMA	MAA	MES	40	40	20	2.5	0.7	1.4	0.8	0.9
	DMAEMA	MAA	MES	40	50	10	4.7	-0.5	-0.5	0.3	1.0
	HEMA-10; MAEMA	MAA	MES	10; 30	40	20	5.7	5.8	0.0	0.3	0.8
	DMAEMA	MAA	MAPTAC	20	66.6	13.3	2.2	8.8	0.1	0.3	0.4
	DMAEMA	MAA	MAPTAC	40	50	10	4.5	8.3	2.3	0.7	1.4


CLAIMS:

1. A composition for washing and/or rinsing of laundry, the composition comprising one or more surfactants suitable for use in laundry wash and/or rinsing products and a polymer which is a co-polymer formed of:-


- (a) one or more anionic monomer units;
- (b) one or more cationic monomer units; and
- (c) optionally, one or more uncharged monomer units;

wherein, the number ratio of the total of all negative charges on the anionic monomer unit(s) to the total of all positive charges on the cationic monomer unit(s) is from 10 : 1 to 3 : 1, especially from 17 : 3 to 3 : 1.

2. A composition according to claim 1, wherein the anionic monomer unit(s) is/are selected from those of formula (A)



- wherein at least two of $\text{Q}^1\text{-Q}^4$ are independently selected from hydrogen and methyl;
- either one or two of $\text{Q}^1\text{-Q}^4$ are independently selected from anionic groups, preferably of formula:



wherein either or both of Q^5 and Q^6 is/are absent, Q^5 otherwise representing -Ph-, -CO-, -CH₂=CH₂, -CONH- or -CO-O- and Q^6 otherwise representing a C₁₋₄ alkylene linkage, one or more of the hydrogen atoms of which is independently optionally substituted by an -OH group or a group -Y;

5

Y is selected from groups of formula -CO₂H, -SO₃H, -OSO₃H, -PO₄H, -PO₃H, -OPO₃H₂ and -OPO₃H₃;

and in the case where two only of Q^1 - Q^4 are independently hydrogen or methyl and only one of Q^1 - Q^4 is - Q^5 - Q^6 -Y, then the remaining group of Q^1 - Q^4 can be any other compatible uncharged group, for example aliphatic, aromatic or mixed aliphatic-aromatic groups having from 2 to 20 carbon atoms (optionally also containing one or more heteroatoms) such as C₂₋₂₀ alkyl groups, C₅₋₁₂ cycloalkyl groups, C₅₋₉ aryl groups, C₁₋₈ alkyl-C₅₋₉ aryl groups, any cycloalkyl or aryl group optionally containing one or two heteroatoms independently selected from nitrogen, oxygen and sulphur.

3. A composition according to either preceding claim, wherein the anionic monomer unit(s) is/are independently selected from monomers having at least one anionic group selected from -CO₂H, -SO₃H, -OSO₃H, -CH₂OSO₃H, -CH=CHSO₃H and groups of formula -(CO)_p-CH₂-CQ⁷Q⁸CO₂H, -PO₄H, -PO₃H, -OPO₃H₂, -OPO₃H₃, wherein p is 0 or 1, Q⁷ is selected from H and OH and Q⁸ is selected from H and CO₂H; and salts thereof.

25


4. A composition according to any preceding claim, wherein the anionic monomer units is/are selected from acrylic acid, methacrylic acid, α -ethacrylic acid, β , β -dimethylacrylic acid, methylenemalononic acid, vinylacetic acid, allylacetic acid, ethylideneacetic acid, propylideneacetic acid, crotonic acid, maleic acid or anhydride, fumaric acid, itaconic acid, citraconic acid, mesaconic acid, N-(methacryloyl)alanine, N-(acryloyl)-hydroxyglycine, sulphopropyl acrylate, sulphoethyl acrylate, sulphoethyl

30

methacrylate, styrenesulphonic acid, vinylsulphonic acid, 2-sulphoethyl methacrylate, sodium allyloxy hydroxypropyl sulphonate, vinylphosphonic acid, phosphoethyl acrylate, phosphonoethyl acrylate, phosphopropyl acrylate, phosphonopropyl acrylate, phosphoethyl methacrylate, phosphonoethyl methacrylate, phosphopropyl methacrylate, phosphonopropyl methacrylate, ethyleneglycol methacrylate phosphate, sulphate of alkoxylate (meth)acrylate, and the salts thereof.


5. A composition according to any preceding claim, wherein the cationic monomer unit(s) is/are independently selected from groups having one or more quaternary nitrogen atoms.

6. A composition according to any preceding claim, wherein the cationic monomer unit(s) is/are independently selected from one or more units derived from compounds of formulae (I) to (III):-




in which:

- R^1 is a hydrogen atom or a methyl group, preferably a methyl group;
- R^2 , R^3 and R^4 are linear or branched $\text{C}_1 - \text{C}_6$ alkyl groups;
- n is from 1 to 4, in particular the number 3;
- Z^1 is a group $-\text{C}(\text{O})\text{O}$, $-\text{C}(\text{O})\text{NH}$ - or $-\text{O}$ -; and
- X^- is a counterion compatible with the water-soluble nature of the polymer;



in which:

- R^5 and R^8 are, independently hydrogen, or a linear or branched $\text{C}_1 - \text{C}_6$ alkyl group;
- R^6 and R^7 are independently represent alkyl, hydroxyalkyl or aminoalkyl group in which the alkyl group is a linear or branched $\text{C}_1 - \text{C}_6$ chain, preferably a methyl group;
- m and p are independently from 1 to 3; and
- X^- is as defined in formula (I); and



in which:

- R^9 is hydrogen, methyl or ethyl;
- $\text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}$ and R^{14} independently selected from groups as defined for R^6 and R^7 in formula(II);
- q is from 0 to 10, preferably from 0 to 2;
- r is from 0 to 6, preferably from 1 to 6, more preferably from 2 to 4;
- Z^1 is as defined in formula (I);
- Z^2 represents a $(\text{CH}_2)_s$ group, s being from 1 to 6, preferably from 2 to 4;
- Z^3 is a linear or branched $\text{C}_2 - \text{C}_{12}$, advantageously $\text{C}_3 - \text{C}_6$, polymethylene chain optionally interrupted by one or more heteroatoms or heterogroups, in particular O or NH, and optionally substituted by one or more hydroxyl or amino groups, preferably hydroxyl groups; and

- X^+ , is as defined in formula (I);
and from those having a cyclic moiety with an N^+ atom.


- 5 7. A composition according to any preceding claim, comprising one or more
uncharged monomer units in a mole ratio of the uncharged units relative to the total of
all anionic monomer units plus the total of all cationic monomer units of from 1 : 1 to
1 : 99, preferably from 3 : 2 to 1 : 9.

10

8. A composition according to any preceding claim, wherein the uncharged
monomer unit(s) is/are selected from

- 15 (i) hydrophilic neutral monomers such as (meth)acrylamide and their N-
monosubstituted or N,N-disubstituted versions (such as N-isopropylacrylamide,
N-butylacrylamide and N,N-dimethylacrylamide), vinyl formamide, vinyl
pyrrolidone, alkoxylated (meth)acrylate, such as hydroxyethyl(meth)acrylate,
hydroxypropyl(meth)acrylate, and their higher ethoxylated or propoxylated
versions such as behenyl polyethoxy methacrylate of formula (V):

20



wherein R^{15} is hydrogen, or methyl and R^{16} is hydrogen, methyl or ethyl, and X is from
1 to 150;

25

- (ii) hydrophobic neutral monomers such as vinyl acetate and its higher homologs,
alkyl(meth)acrylates (e.g. methyl methacrylate, butyl acrylate and ethyl

acrylate), styrene and its derivatives, methyl vinyl ether, Sipomer WAM and WAM II from Rhodia, glycidyl methacrylate; and

5 (iii) hydrophilic neutral monomers with potentially cationic functional groups.

9. A laundry wash composition according to any preceding claim, wherein the surfactant(s) is/are selected from one or more anionic and/or nonionic surfactants.

10 10. A laundry wash composition according to any preceding claim, wherein the composition, when diluted in wash liquor, gives a pH of the wash liquor from 7 to 11, preferably from 7 to 10.5.

11. A laundry rinse composition according to any of claim 1-6, wherein the
15 surfactant(s) is are selected from one or more cationic rinse conditioner surfactants.

12. Use of a polymer for a method effecting soil-release of a laundry item, the method comprising contacting the laundry item with the said polymer, the polymer being a copolymer formed of:-
20

- (a) one or more anionic monomer units;
- (b) one or more cationic monomer units; and
- 25 (c) optionally, one or more uncharged monomer units;

wherein, the number ratio of the total of all negative charges on the anionic monomer unit(s) to the total of all positive charges on the cationic monomer unit(s) is from 10 : 1 to 3 : 1, especially from 17 : 3 to 3 : 1.

INTERNATIONAL SEARCH REPORT

national Application No

PLT/EP 01/00409

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11D3/37 C11D1/62 C11D3/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 196 49 288 A (HENKEL KGAA) 4 June 1998 (1998-06-04) page 4, line 33 - line 36; claims 1-7,13,14; examples; table 1 page 8, line 25 - line 31 & WO 98 23714 A (HENKEL) 4 June 1998 (1998-06-04) cited in the application	1-10,12
X	WO 98 16538 A (UNILEVER PLC ;GRAINGER DAVID STEPHEN (GB); UNILEVER NV (NL); KHAN) 23 April 1998 (1998-04-23) claim 1; examples 50D,53	1-6,9
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

25 June 2001

Date of mailing of the international search report

17/07/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Loiselet-Taisne, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/00409

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 198437 Derwent Publications Ltd., London, GB; Class A97, AN 1984-228141 XP002141108 & JP 59 135293 A (KAO CORP), 3 August 1984 (1984-08-03) abstract	1-11
X	US 5 413 731 A (ADLER DAVID E ET AL) 9 May 1995 (1995-05-09) column 10, line 42 - line 50; claims 1-16; table 1	1-9
A	US 5 783 533 A (KENSICHER YVES ET AL) 21 July 1998 (1998-07-21) trial 2, 5-8, 22 column 21, line 41 - line 44; claims 1,4,8,12,16,17; example 7; table 5	1-4
A	GB 2 104 091 A (KAO CORP) 2 March 1983 (1983-03-02) cited in the application synthesis ex. 1, 2 page 3, line 42 - line 48; table 1	1-3

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/EP 01/00409

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 19649288	A	04-06-1998	WO 9823714 A EP 0944701 A	04-06-1998 29-09-1999
WO 9816538	A	23-04-1998	AU 723907 B AU 5051098 A BR 9711906 A EP 0934328 A TR 9900801 T ZA 9709231 A	07-09-2000 11-05-1998 24-08-1999 11-08-1999 21-07-1999 15-04-1999
JP 59135293	A	03-08-1984	JP 1792760 C JP 5000439 B	14-10-1993 05-01-1993
US 5413731	A	09-05-1995	US 5308532 A AT 169331 T AU 3388293 A BR 9301108 A CA 2090932 A DE 69320082 D DE 69320082 T EP 0560519 A JP 6009726 A KR 252576 B MX 9301285 A	03-05-1994 15-08-1998 16-09-1993 14-09-1993 11-09-1993 10-09-1998 06-05-1999 15-09-1993 18-01-1994 15-04-2000 01-09-1993
US 5783533	A	21-07-1998	FR 2732031 A CA 2169894 A EP 0733700 A	27-09-1996 24-09-1996 25-09-1996
GB 2104091	A	02-03-1983	JP 1055320 B JP 1569529 C JP 58013700 A HK 46788 A MY 54987 A SG 63187 G	24-11-1989 10-07-1990 26-01-1983 01-07-1988 31-12-1987 19-02-1988